

Cluster Aspects of ^{252}Cf Spontaneous Fission

R. Donangelo¹, J.O. Rasmussen, M.A. Stoyer² and J.H. Hamilton³

Permanent Addresses:

¹ *Instituto de Física, Universidade Federal do Rio de Janeiro,
21945-970 Rio de Janeiro, Brazil*

² *Lawrence Livermore National Laboratory, Livermore, CA 94550*

³ *Department of Physics and Astronomy, Vanderbilt University
Nashville, Tennessee 37235, USA*

Based on gamma coincidence data obtained with Gammasphere yields of specific two-fragment combinations can be directly determined in a two-dimensional yield matrix. The largest matrix of ^{252}Cf spontaneous fission product yield data comes from the Mo-Ba split. These yields, which are shown in Ref. [1], show that this split is unique in that it has an unusual component of 8-10 neutron-emission at a level of about 7% of the total integrated yields of Mo-Ba.

The idea of two co-existing fission modes is not new. Hulet *et al* [2] suggested the occurrence of two different fission processes leading to two modes of fission. What is new in the spontaneous fission of ^{252}Cf is that the low $\langle TKE \rangle$ mode manifests itself through the same primary mass split $^{106}\text{Mo}-^{146}\text{Ba}$, which is one of the most abundant primary pairs emerging in the “normal” mode.

In order to extract the essential physical ingredients characterizing the data we performed a study which considered a minimum number of adjustable parameters and made use of whatever constraints on parameters available from other fission data. The details of this study are given in Ref. [3], and its main result is that it associates the low $\langle TKE \rangle$ (i.e. highly excited fragments) mode to a cluster structure of the Ba fragment at scission, in which a ^{14}C forms a neck between the Mo fragment and a ^{132}Sn cluster. After scission, the C and Sn clusters recombine, forming a highly excited Ba fragment. This process accounts for the apparent unequal partition of excitation energy between the Ba and the Mo fragments, and furthermore appears to be in agree-

ment with the recent work of Pyatkov *et al.* [4], which show more than one valley as scission is approached, some of which leading to shapes of the fissioning system close to what we would expect of a quasi-ternary mode as the one described above.

References

- [1] J.H. Hamilton *et al.*, Prog. Particle and Nucl. Phys. 35(1995)635.
- [2] E.K. Hulet *et al.* Phys. Rev. Lett. 56(1986)313.
- [3] R. Donangelo *et al.*, To be published in the Int. J. of Mod. Phys. E (1998).
- [4] Yu.V. Pyatkov *et al.*, “Cluster degrees of freedom in fission of Californium nuclei” in Proc. Int. Conf. on Dynamical Aspects of Nuclear Fission – Casta-Papiernicka, Slovakia, 1996, eds. J. Kliman and B. Pustylnik JINR, Dubna 1996, p. 322.